Chapter 6 - Virology

- Topics
 - -Structure
 - -Classification
 - Multiplication
 - -Cultivation and replication
 - -Nonviral infectious agent
 - -Teratogenic/Oncogenic

- Viruses have a **host range**. That is, viruses infect specific cells or tissues of specific hosts, or specific bacteria, or specific plants.
- Viral specificity refers to the specific kinds of cells a virus can infect. It is regulated by the specificities of attachment, penetration and replication of the virus (Receptors)

Properties of viruses

- Viruses are not cells, do not have nuclei or mitochondria or ribosomes or other cellular components.
- Viruses replicate or multiply. Viruses do not grow.
- Viruses replicate or multiply only within living cells.
- Viruses are obligate intracellular parasites.
- The term virus was coined by Pasteur, and is from the Latin word for **poison**.

Components of viruses -


A virion is an infectious virus particle - not all virus particles are infectious


Viruses are composed of a nucleic acid, RNA or DNA - never both.

- All viruses have a **protein coat** (capside) or shell that surrounds and protects the nucleic acid core.
- Some viruses have a lipid envelope or membrane surrounding a nucleocapsid core. The source of the envelope is from the membranes of the host cell.
- Some viruses package enzymes e.g. RNAdependent-RNA polymerase or other enzymes some do not package enzymes

Structure

- Size and morphology
- Capsid
- Envelope
- Complex
- Nucleic acid





- Protective outer shell that surrounds viral nucleic acid
- Composed of capsomer subunits - collectively protect the nucleic acid from the environment
- Capsid spikes used for binding to cell surface proteins

Envelope


- · Lipid and proteins basically a modified version of our membranes
- Envelope spikes bind to cell surface proteins
- During release of animal viruses, a part of the host membrane is taken


Nucleic acid


- Viruses contain either DNA or RNA
- Possess only the genes to invade and regulate the metabolic activity of host cells
- Ex. Hepatitis B (4 genes) and herpesviruses (100 genes)
- No viral metabolic genes, as the virus uses the host's metabolic resources


Bacteriophages


- Bacteriophage
 - -Polyhedral head
 - -Helical tail
 - -Fibers for attachment
- Are considered either LYTIC or TEMPERATE
- · Are often associated with virulence genes in bacteria
 - EX. diphtheria toxin in *Clostridium* diphtheriae - also Botulinum toxin from C. botulinum















Temperate phages can cause disease

For example, *Corynebacterium diphtheriae* and *Clostridium botulinum* contain **prophages** that have genes which encode for toxins


Without these prophages, they **DO NOT** produce the toxin – without toxin, no disease

Thus, they are examples of bacteria and viruses interacting to cause medically relevant disease



Classification


- · Host Range
- · Envelope or Naked virus
- · Type of disease
- Baltimore Classification of Viruses (4 major groups)
 - + vs sense





General Steps in Viral Multiplication


- Adsorption
- Penetration
- Uncoating
- Synthesis
- Assembly
- Release

Viruses recognize specific receptors
Figure 6.12 and then the virus penetrates the cell Figure 6.13












Animal Virus Replication


General Steps in Viral Multiplication


- Adsorption
- Penetration
- Uncoating
- Synthesis
- Assembly
- Release
















Replication of RNA viruses Insight 6.2 - Note differences between + sense, - sense and retrovirus replication

Also, be able to compare -contrast phage and animal virus replication

Cytopathic effects

- Damage to the host cell due to a viral infection
 - -Transformation





Cultivation and Replication


- · In vivo methods
 - -Laboratory animals
 - -Embryonic bird tissues
- · In vitro methods
 - -Cell or tissue culture


Cultivation of animal viruses -

It is possible to study viruses in animals, but due to the complexity of the animal, expense of animals and the political environment relative to the use of animals, alternatives have been developed.

Chick embryos in eggs - influenza

Persistent infection - transformation/cancer






Noncellular Infectious Agents

Prions

Prions – Stanley Pruisner – Nobel Prize in Medicine: 1978

Prions are proteinaceous infectious agents - Prions do not have nucleic acid.

Prions are considered to be the causative agents of Creutzfeld Jakob disease, scrapie, bovine spongiform encephalopathy (BSE) and kuru.



Oncogenic potential of viruses -

Cancer is a set of diseases known to disturb the normal functioning and properties of cells.

Tumors may be malignant or benign - malignant tumors spread by metastasis.

Peyton Rous in 1911 discovered that a filterable agent could transmit a sarcoma (a type of cancer) in chickens - Rous sarcoma virus - the first retrovirus described.

At least six viruses have been found to cause human cancer - Epstein-Barr virus, hepatitis B virus, hepatitis C virus, human papilloma virus (HPV-8, HPV-16), HTLV-I (adult T-cell leukemia and lymphoma), HTLV-II (hairy cell leukemia).

Oncogenes are normal cellular regulatory genes. When modified, these genes code for gene products that disturb the normal regulatory patterns of cells and can result in a loss of the normal properties of cell growth and division resulting in "cancer".

Viral oncogenes are found usually in retroviruses.

Viral oncogenes are viral homologs of the cellular oncogenes.

Viral oncogenes can disturb normal regulatory properties by certain mechanisms.

