

# Net primary productivity and rain-use efficiency as affected by warming, altered precipitation, and clipping in a mixed-grass prairie

XIA XU\*, REBECCA A. SHERRY\*, SHULI NIU\*†, DEJUN LI\* and YIQI LUO\*

\*Department of Microbiology and Plant Biology, University of Oklahoma, 101 David L. Boren Blvd, Norman, OK 73019, USA,

†Key Laboratory of Ecosystem Network Observation and Modeling, Institute of Geographic Sciences and Natural Resources Research, Chinese Academy of Sciences, Beijing 100101, China

## Abstract

Grassland productivity in response to climate change and land use is a global concern. In order to explore the effects of climate change and land use on net primary productivity (NPP), NPP partitioning [ $f_{\text{BNPP}}$ , defined as the fraction of belowground NPP (BNPP) to NPP], and rain-use efficiency (RUE) of NPP, we conducted a field experiment with warming (+3 °C), altered precipitation (double and half), and annual clipping in a mixed-grass prairie in Oklahoma, USA since July, 2009. Across the years, warming significantly increased BNPP,  $f_{\text{BNPP}}$ , and  $\text{RUE}_{\text{BNPP}}$  by an average of 11.6%, 2.8%, and 6.6%, respectively. This indicates that BNPP was more sensitive to warming than aboveground NPP (ANPP) since warming did not change ANPP and  $\text{RUE}_{\text{ANPP}}$  much. Double precipitation stimulated ANPP, BNPP, and NPP but suppressed  $\text{RUE}_{\text{ANPP}}$ ,  $\text{RUE}_{\text{BNPP}}$ , and  $\text{RUE}_{\text{NPP}}$  while half precipitation decreased ANPP, BNPP, and NPP but increased  $\text{RUE}_{\text{ANPP}}$ ,  $\text{RUE}_{\text{BNPP}}$ , and  $\text{RUE}_{\text{NPP}}$ . Clipping interacted with altered precipitation in impacting  $\text{RUE}_{\text{ANPP}}$ ,  $\text{RUE}_{\text{BNPP}}$ , and  $\text{RUE}_{\text{NPP}}$ , suggesting land use could confound the effects of precipitation changes on ecosystem processes. Soil moisture was found to be a main factor in regulating variation in ANPP, BNPP, and NPP while soil temperature was the dominant factor influencing  $f_{\text{BNPP}}$ . These findings suggest that BNPP is critical point to future research. Additionally, results from single-factor manipulative experiments should be treated with caution due to the non-additive interactive effects of warming with altered precipitation and land use (clipping).

**Keywords:** above- and belowground net primary productivity, climate change, clipping, mixed-grass prairie, net primary productivity partitioning, rain-use efficiency

Received 24 October 2012; revised version received 20 April 2013 and accepted 27 April 2013

## Introduction

In response to rising concentrations of atmospheric greenhouse gases, global mean temperature has increased 0.74 °C since 1850 and is predicted to increase another 1.8–4.0 °C by the end of this century (IPCC, 2007). Additionally, precipitation is anticipated to vary greatly under global climate change, with increasing intra- and inter-annual variability (IPCC, 2007). Net primary productivity (NPP), the primary driver of global carbon (C) cycling, is strongly influenced by climatic variables, such as temperature and precipitation (e.g., Rustad *et al.*, 2001; Sherry *et al.*, 2008; Bloor & Bardgett, 2012; Hoeppepner & Dukes, 2012). Both empirical and modeling studies have suggested that climatic changes could alter the functioning of terrestrial ecosystems via changes in NPP (e.g., Norby & Luo, 2004; Luo *et al.*, 2009). However,

responses of NPP to climate warming, altered precipitation, and their interactions are less clear (Rustad *et al.*, 2001), especially belowground NPP (BNPP). Understanding variation in NPP is critical to discovering the mechanisms of the response of ecosystem functions to ongoing climate change (Ni, 2004; Xu *et al.*, 2012).

Evidence that warming and altered precipitation are influencing ecosystem processes is growing rapidly (e.g., Ni, 2004; Luo *et al.*, 2009; Wu *et al.*, 2011; Polley *et al.*, 2012; Xu *et al.*, 2012). Manipulative experiments in a range of vegetation types have shown that warming may have a variety of impacts on NPP, either stimulating plant productivity (Rustad *et al.*, 2001) or retarding it (Rustad *et al.*, 2001; Wu *et al.*, 2011). Studies examining the effects of altered precipitation on plant productivity have shown that positive correlations typically exist between plant productivity and annual precipitation in grassland ecosystems (Sala *et al.*, 1988; Knapp & Smith, 2001; Dukes *et al.*, 2005; Chimner *et al.*, 2010). However, it is less clear whether the positive relationship holds true under

Correspondence: Dr Xia Xu, tel. +405 325 6519, fax +405 325 7619, e-mail: xuxia.1982@yahoo.com

extremely altered precipitation, such as double and half precipitation, which has been recognized as a critical manifestation of climate change (Meehl *et al.*, 2007). Moreover, the interactions of warming and altered precipitation may shape the NPP in ways that are hard to predict from measuring responses of an ecosystem to single climate change factors (Dukes *et al.*, 2005; Williams *et al.*, 2007). The variability and unpredictability of NPP under climatic change may also result from our limited understanding of BNPP that represents more than one-half of NPP in grasslands (Hui & Jackson, 2005). In comparison to reasonably well-understood aboveground NPP (ANPP), experimental estimates of BNPP are less and only a few studies have investigated the relation of BNPP with climatic factors (Gill *et al.*, 2002; Ni, 2004; Dukes *et al.*, 2005; Xu *et al.*, 2012). The lack of NPP data, especially the magnitude and direction of its response to current climate change, largely inhibits the progress in validating global NPP models and projecting future climate change (Cramer *et al.*, 1999; Hui & Jackson, 2005).

Net primary productivity partitioning, commonly defined as  $f_{\text{BNPP}} = \text{BNPP} / (\text{ANPP} + \text{BNPP})$ , is a critical constraint for the calibration and testing of dynamic C-cycling models (Ågren & Franklin, 2003; Hui & Jackson, 2005). From a physiological perspective,  $f_{\text{BNPP}}$  is sensitive to environmental changes (Bloom *et al.*, 1985; Chapin *et al.*, 1987). For example, drought and warming appear to favor allocation of C from leaves to roots (Poorter & Nagel, 2000; Xu *et al.*, 2012). However, it is generally assumed that  $f_{\text{BNPP}}$  is constant in a majority of global terrestrial models (e.g., Friedlingstein *et al.*, 1999; Wullschlegel *et al.*, 2001). The partitioning of NPP is less well-understood and remains under debate (Enquist & Niklas, 2002; Shipley & Meziane, 2002), especially in relation to climatic changes, leading to tremendous variation for short-term forecasts and long-term projections of global NPP (Friedlingstein *et al.*, 2006; Sitch *et al.*, 2008). Thus, results from field manipulative experiments are fundamentally important to understanding and quantifying  $f_{\text{BNPP}}$  and its relationships with climatic factors.

Efficient rain use is generally one of the key components of adaptability for natural vegetation (Brueck *et al.*, 2010). Rain-use efficiency (RUE), the ratio of NPP to precipitation, provides us a useful index for evaluating the responses of NPP to precipitation changes (Knapp *et al.*, 2001; Brueck *et al.*, 2010) and is an important constraint for simulating plant productivity in models (Roupsard *et al.*, 2009). However, how RUE will vary under warming and altered precipitation is not clear. In grasslands, RUE either increases (Xu *et al.*, 2012) or decreases (De Boeck *et al.*, 2006) under

warming, although it is predicted to decrease in a modeling study (Bell *et al.*, 2010). Similarly, increases (Epstein *et al.*, 1996; Hooper & Johnson, 1999), no changes (Lauenroth *et al.*, 2000), and decreases (Huxman *et al.*, 2004; Bai *et al.*, 2008) in RUE have been reported with increasing precipitation amount. Therefore, greater understanding of how RUE responds to climatic changes is critical to accurately forecast terrestrial C-cycle response and feedback to climate change.

Land-use practices, such as mowing for hay, may further confound the responses of NPP,  $f_{\text{BNPP}}$ , and RUE of NPP to climate change (McNaughton *et al.*, 1998; Gao *et al.*, 2008) and impact ecosystem C balance (Luo *et al.*, 2009). Hay harvest is a widely practiced land use in the southern Great Plains of the US. Hay production occupies 3.25 million acres in Oklahoma, nearly as much as wheat (USDA, National Agricultural Statistic Service). Clipping to mimic hay harvest has considerable impacts on ANPP because it directly takes aboveground biomass away, lowers the amount of litter on the ground, and indirectly impacts plant growth in the next growing season. On the other hand, clipping influences BNPP and  $f_{\text{BNPP}}$  by breaking down inherent allocation rules due to lesser demand for water and nutrients from aboveground plant components (Xu *et al.*, 2012). Additionally, clipping may affect NPP and RUE by stimulating evapotranspiration, reducing soil moisture, and exacerbating water stress (Niu *et al.*, 2008; Xu *et al.*, 2012). Although NPP has been reported to be affected by temperature, water availability, nitrogen availability, and grazing (Ni, 2004; Sherry *et al.*, 2008; Gao *et al.*, 2011; Li *et al.*, 2011; Xu *et al.*, 2012), experimental estimates of NPP in response to clipping or haying are still few, making it difficult to predict the potential ecosystem-level responses to land-use practices under climate change in grasslands.

Grassland ecosystems, accounting for ca. 54% of the conterminous US (US Department of Agriculture, 1972), play an essential role in global C cycling. Grasslands are ideal for addressing the potential responses of NPP and its RUE to climate change and land use (Hui & Jackson, 2005; Xu *et al.*, 2012) because of (i) their rapid responses to climate change and land-use practices (Sherry *et al.*, 2008; Luo *et al.*, 2009); and (ii) the relative ease of manipulating temperature, precipitation, and land-use practices, of measuring belowground biomass (Gill *et al.*, 2002). Specifically, we aimed to: (i) examine the effects of warming, altered precipitation, clipping, and their interactions on NPP,  $f_{\text{BNPP}}$ , and RUE of NPP; and (ii) explore the relationships of NPP and  $f_{\text{BNPP}}$  with climatic factors and species composition (expressed by the ratio of  $C_3$  contribution to ANPP) in a mixed-grass prairie in the southern Great Plains of the USA.

## Materials and methods

### Experimental site and design

The experimental site is located on the Kessler Atmospheric and Ecological Field Station (KAEFS) in McClain County, Oklahoma, USA (ca. 34°59'N, 97°31'W), about 40 km southwest of the Norman campus of the University of Oklahoma, USA. KAEFS is located in the Central Redbed Plains of Oklahoma within the mosaic ecotone from tall-grass prairie with woodlands along creeks to mixed-grass prairie in the west (Tarr *et al.*, 1980). The site is on an old-field prairie abandoned from field cropping 40 years ago with light grazing until 5 years ago. The mixed-grass prairie is dominated by C<sub>3</sub> forbs (*Ambrosia trifida*, *Solanum carolinense*, and *Euphorbia dentata*) and C<sub>4</sub> grasses (*Tridens flavus*, *Sporobolus compositus*, and *Sorghum halapense*). Mean annual temperature is 16.3 °C and mean annual precipitation is 914 mm (Oklahoma Climatological Survey, Norman, OK, USA). The soil is part of the Nash-Lucien complex with neutral pH, high available water holding capacity (around 37%), and a deep (ca. 70 cm), moderately penetrable root zone.

This experiment was established in July of 2009 and manipulates temperature and precipitation, within which is nested a clipping factor. Each treatment is randomly repeated four times for a total 24 plots of 2.5 m × 3.5 m. We utilize infrared heaters to achieve the whole ecosystem warming by ca. 3 °C. In each warmed plot, two infrared heaters (165 cm × 15 cm; Kalglo Electronics, Bethlehem, PA, USA) are suspended ca. 1.5 m above the ground to warm the area of 2.5 m × 1.75 m. The control plot has two 'dummy' heaters with same dimensions as the infrared heaters suspended at a similar height to mimic the shading effects of the heaters. Temperature increments generated by the infrared heaters are found relatively even over the entire area of the plots and similar at different soil depths (Wan *et al.*, 2002). The distance between each two adjacent plots is at least 5 m from centers to avoid heating of the control plots.

We use a rainfall-collection-redistribution device as described by Zhou *et al.* (2006) with a same area of the plot to double precipitation and a rainout-shelter as described by Yahdjian & Sala (2002) to halve precipitation. The rainout-shelter design is a fixed-location shelter with a roof consisting of bands of transparent acrylic that block different amounts of rainfall while minimally affecting other environmental variables. To minimize disturbance, we inserted fiberglass sheets into the ground to a depth of 120 cm around each plot as in the Jasper Ridge Global Change Experiment (Zavaleta *et al.*, 2003) to cut off lateral movement of soil water. We set the rain shelters above the heaters at an angle of 20° as in the original design by Yahdjian & Sala (2002). The lower side of the rainout-shelters tilts toward the prevailing upwind direction. In the control plots, we install 'dummy' frames of the rain shelters for consistency of experimental conditions.

Each 2.5 m × 3.5 m plot is divided into two 2.5 m × 1.75 m subplots. Plants in the southern 2.5 m × 1.75 m subplots are clipped at a height of 10 cm above the ground once a year to mimic the land-use practice of mowing for hay while the northern subplots are unclipped. Similar to the hay production,

clipped materials are taken away and not returned back to the plots. Thus, this experiment has twelve treatments, six for the unclipped group and the other six for the clipped group. The six treatments are control (ambient) temperature and control precipitation (CC), control temperature and double precipitation (CD), control temperature and half precipitation (CH), warming and control precipitation (WC), warming and double precipitation (WD), and warming and half precipitation (WH).

### Soil temperature and moisture, and precipitation measurements

Soil temperature, at the depth of 7.5 cm in the center of one clipped and one unclipped subplot, was measured every 15 min using thermocouples (T-type; Campbell Science Inst., Logan, UT, USA) wired to a Campbell Scientific CR10x datalogger (Campbell Scientific). Volumetric soil water content (%V) in the top 12 cm was measured once or twice a month using portable Time Domain Reflectometry (TDR) equipment (Soil Moisture Equipment Corp., Santa Barbara, CA, USA). Three measurements of soil moisture were made in every subplot each time and the average values were used in analysis. Precipitation data were obtained online from an Oklahoma Mesonet Station (Washington Station) located ca. 50 m away from our experimental site. The amount of actual precipitation received by the plots under different precipitation treatments was measured using surface gauge measurement (Groisman & Legates, 1994; New *et al.*, 2001).

### NPP measurement and estimation of $f_{\text{BNPP}}$ and RUE of NPP

In 2010 and 2011, ANPP, separated into C<sub>3</sub> and C<sub>4</sub> species, was directly measured by annual clipping at peak biomass (usually August) in the southern 2.5 m × 1.75 m clipping subplots and indirectly estimated by pin-contact method (Frank & McNaughton, 1990) in the unclipped subplots. Detailed description of biomass estimation is provided by Sherry *et al.* (2008). The root ingrowth-core method (Gao *et al.*, 2008; Xu *et al.*, 2012) was applied to estimate BNPP. Soil cores (5.2 cm in diameter, 90 cm in length) were taken once a year in October at an angle of 90° from the same spots in one unclipped and one clipped subplots of each plot every year. Holes were immediately refilled with sieved root-free soils originating from the same depth (4 depths in total: 0–15, 15–30, 30–60, 60–90 cm) outside of the plots. Soil filled into the holes was compressed to a density comparable to the bulk soil. Separated soil cores were put into plastic bags, transported in several coolers to the Ecolab at the University of Oklahoma, Norman and stored at –30 °C before analyzing. Root samples were carefully washed by wet sieving (0.5 mm) under gently flowing water to remove attached soil and dark brown/black debris, oven-dried at 70 °C for 48 h and weighted to calculate BNPP.  $f_{\text{BNPP}}$  is defined as  $f_{\text{BNPP}} = \text{BNPP}/(\text{ANPP} + \text{BNPP})$  according to Hui & Jackson (2005). Rain-use efficiency (RUE) of ANPP, BNPP, and NPP is calculated as ANPP, BNPP, and NPP, respectively, divided by the amount of actual precipitation received by specific plots under different treatments. RUE

of different components of NPP was estimated to compare the sensitivity between ANPP and BNPP to climate change and land use.

### Statistical analysis

Repeated measures split-plot analysis of variance (ANOVA) was used to examine the main and interactive effects of experimental warming (whole-plot factor), altered precipitation (whole-plot factor), clipping (subplot factor), and year on ANPP, BNPP, NPP,  $f_{\text{BNPP}}$ ,  $\text{RUE}_{\text{ANPP}}$ ,  $\text{RUE}_{\text{BNPP}}$ ,  $\text{RUE}_{\text{NPP}}$ , and soil moisture and temperature. Within each year, one-way ANOVA was performed to analyze the differences in ANPP, BNPP, NPP,  $f_{\text{BNPP}}$ ,  $\text{RUE}_{\text{ANPP}}$ ,  $\text{RUE}_{\text{BNPP}}$ , and  $\text{RUE}_{\text{NPP}}$  among the twelve treatments (Duncan's Test) and to test the separate main treatment effects on the parameters mentioned above (Tukey's Test). Linear regression analyses were performed to evaluate the relationships of ANPP, BNPP, NPP, and  $f_{\text{BNPP}}$  with climatic factors and species composition (expressed by the ratio of  $C_3$  contribution to ANPP). All statistical analyses were conducted using SPSS 16.0 for windows (SPSS Inc., Chicago, IL, USA).

## Results

### Microclimate

Comparing with the average precipitation of 837 mm from 1994 to 2011, annual precipitation in 2010 and 2011 was 906 mm and 549 mm, respectively (Table 1; Fig. 1). The amount of precipitation received by the control, double, and half precipitation treatment plots differed significantly (all  $P < 0.01$ ). According to our measurements, plots under double and half precipitation treatments received an average of 195.8% and 70.1% of the ambient precipitation, respectively (Fig. 1c). Overall, warming, altered precipitation, and clipping significantly affected both soil moisture and soil temperature (all  $P < 0.01$ , Table 2). Averaged across years, volumetric soil moisture was lowered by 1.9%, 0.3%, and 0.8% by the treatments of warming, half precipitation, and clipping, respectively, and increased by an average of 1.1% under double precipitation (Table 1). Additionally, warming, double precipitation, half precipitation, and clipping increased soil temperature by an average of 3.1, 0.1, 0.4, and 0.5 °C, respectively, across the years (Table 1).


### Treatment effects on NPP and $f_{\text{BNPP}}$

Ecosystem productivity varied greatly from 2010 to 2011. Overall, the productivity in 2011 was much smaller than that in 2010 (Table 2; Fig. 2a–c). ANPP varied from  $184.0 \pm 23.1 \text{ g m}^{-2}$  in 2011 under the treatment of warming plus half precipitation in the clipped subplots to  $561.9 \pm 39.7 \text{ g m}^{-2}$  in 2010 under the control (Fig. 2a). The main effects of warming increased BNPP and NPP by an average of 11.6% ( $P < 0.01$ ) and 9.2%

**Table 1** Annual precipitation (PPT), mean soil moisture ( $W_{\text{soil}}$ ), and mean soil temperature ( $T_{\text{soil}}$ ) under 12 treatments in 2010 and 2011

Year	PPT (mm)	Treatment	Unclipped						Clipped					
			CC	CD	CH	WC	WD	WH	CC	CD	CH	WC	WD	WH
2010	906	$W_{\text{soil}}$ (%)	11.6 ± 0.2	13.0 ± 0.3	11.6 ± 0.6	9.6 ± 0.5	12.1 ± 1.1	8.7 ± 0.6	10.6 ± 0.6	12.3 ± 0.5	11.4 ± 0.3	9.2 ± 0.6	8.8 ± 0.3	7.7 ± 0.6
		$T_{\text{soil}}$ (°C)	16.2 ± 0.0	16.3 ± 0.0	16.7 ± 0.1	18.5 ± 0.0	18.9 ± 0.0	18.6 ± 0.0	16.6 ± 0.0	16.7 ± 0.4	16.9 ± 0.0	19.2 ± 0.0	19.6 ± 0.0	20.0 ± 0.4
2011	549	$W_{\text{soil}}$ (%)	7.6 ± 0.4	9.1 ± 0.2	7.3 ± 0.5	6.3 ± 0.2	7.2 ± 0.2	5.9 ± 0.1	7.0 ± 0.6	7.9 ± 0.6	6.9 ± 0.4	6.0 ± 0.2	6.5 ± 0.3	5.6 ± 0.2
		$T_{\text{soil}}$ (°C)	18.3 ± 0.1	18.1 ± 0.1	18.4 ± 0.1	21.9 ± 0.1	21.7 ± 0.1	21.7 ± 0.1	18.5 ± 0.1	18.1 ± 0.0	18.8 ± 0.1	21.7 ± 0.1	22.0 ± 0.1	22.9 ± 0.1

CC, control temperature plus control PPT; CD, control temperature plus double PPT; CH, control temperature plus half PPT; WC, warmed plus control PPT; WD, warmed plus double PPT; WH, warmed plus half PPT.



**Fig. 1** Daily (a) and monthly (b) distribution of precipitation in 2010 and 2011 and the effectiveness of precipitation treatments (c). Different letters represent statistically significant differences at  $P < 0.05$ . C, control; D, double precipitation; H, half precipitation.

( $P = 0.057$ , Table 2; Fig. 3a), respectively. Altered precipitation significantly influenced ecosystem productivity across the years (all  $P < 0.05$ , Table 2). For example, the main effects of double precipitation increased ANPP, BNPP, and NPP by an average of 3.9%, 3.8%, and 3.9%, respectively (Fig. 3c). In contrast, the main effects of half precipitation decreased ANPP, BNPP, and NPP by an average of 13.5%, 10.9%, and 12.2%, respectively (Fig. 3e). Overall, clipping had no significant impact on ANPP, BNPP, and NPP (all  $P > 0.05$ , Table 2; Fig. 3g).

$f_{\text{BNPP}}$  varied greatly from  $0.5 \pm 0.0$  in 2010 under the control to  $0.6 \pm 0.0$  in 2011 under warming plus

clipping in the clipped subplots (Fig. 2d). Warming significantly increased  $f_{\text{BNPP}}$  across the years ( $P < 0.05$ , Table 2; Fig. 3b). But neither the main effects of altered precipitation nor the impact of clipping was significant on  $f_{\text{BNPP}}$  over the years (all  $P > 0.05$ , Table 2). The main effects of warming and altered precipitation on  $f_{\text{BNPP}}$  varied between 2010 and 2011 ( $P = 0.042$  and  $0.057$ , respectively, Table 2). No interactive effects of warming, altered precipitation, and clipping on  $f_{\text{BNPP}}$  were found (all  $P > 0.05$ , Table 2).


#### Treatment effects on RUE of NPP

The treatments and some of their interactions significantly influenced RUE of NPP across the years (Table 2; Fig. 4). Although warming effects on  $\text{RUE}_{\text{ANPP}}$  and  $\text{RUE}_{\text{NPP}}$  were not significant (all  $P > 0.05$ ), warming increased  $\text{RUE}_{\text{BNPP}}$  by an average of 5.0% across the years ( $P < 0.05$ , Table 2; Fig. 5a). Generally, altered precipitation greatly affected  $\text{RUE}_{\text{ANPP}}$ ,  $\text{RUE}_{\text{BNPP}}$ , and  $\text{RUE}_{\text{NPP}}$  (all  $P < 0.01$ , Table 2; Fig. 4). Double precipitation on average decreased  $\text{RUE}_{\text{ANPP}}$ ,  $\text{RUE}_{\text{BNPP}}$ , and  $\text{RUE}_{\text{NPP}}$  by 24.3%, 26.0%, and 50.4%, respectively, across the years (Fig. 5b). In contrast, the average increases under half precipitation treatment in  $\text{RUE}_{\text{ANPP}}$ ,  $\text{RUE}_{\text{BNPP}}$ , and  $\text{RUE}_{\text{NPP}}$  were 11.6%, 14.3%, and 25.9%, respectively (Fig. 5c). Clipping marginally decreased  $\text{RUE}_{\text{ANPP}}$ ,  $\text{RUE}_{\text{BNPP}}$ , and  $\text{RUE}_{\text{NPP}}$  by an average of 4.5% ( $P = 0.075$ ), 3.1% ( $P = 0.118$ ), and 7.6% ( $P = 0.083$ ), respectively, across the years (Table 2; Fig. 5d). The main treatment effects on  $\text{RUE}_{\text{ANPP}}$ ,

**Table 2** Results of repeated measures split-plot ANOVA for responses of soil temperature (ST) and moisture (SM), ANPP, BNPP, NPP,  $f_{\text{BNPP}}$ , and RUE of ANPP, BNPP, and NPP to warming (W), altered precipitation (PPT), clipping (C), year, and their interactions ( $n = 4$ ).  $P$  values smaller than 0.05 and 0.10 are in bold and italic, respectively

	ST	SM	ANPP	BNPP	NPP	$f_{\text{BNPP}}$	$\text{RUE}_{\text{-ANPP}}$	$\text{RUE}_{\text{-BNPP}}$	$\text{RUE}_{\text{-NPP}}$
Warming (W)	<b>0.001</b>	<b>&lt;0.001</b>	0.217	<b>0.009</b>	<i>0.057</i>	<b>0.020</b>	0.454	<b>0.013</b>	0.114
PPT	<b>0.002</b>	<b>&lt;0.001</b>	<b>0.024</b>	<b>0.010</b>	<b>0.015</b>	0.234	<b>&lt;0.001</b>	<b>&lt;0.001</b>	<b>&lt;0.001</b>
Clipping (C)	<b>0.007</b>	<b>&lt;0.001</b>	0.195	0.157	0.170	0.340	<i>0.075</i>	0.118	<i>0.083</i>
Year (Y)	<b>&lt;0.001</b>	<b>&lt;0.001</b>	<b>&lt;0.001</b>	<b>&lt;0.001</b>	<b>&lt;0.001</b>	<b>&lt;0.001</b>	<b>&lt;0.001</b>	0.493	<b>0.029</b>
W × PPT	<b>&lt;0.001</b>	0.193	0.311	0.211	0.286	0.227	0.435	0.139	0.303
W × C	<b>&lt;0.001</b>	0.516	0.451	0.254	0.348	0.634	0.616	0.369	0.487
W × Y	<b>&lt;0.001</b>	<b>&lt;0.001</b>	0.983	0.422	0.702	<b>0.042</b>	0.953	<i>0.082</i>	0.355
PPT × C	<b>&lt;0.001</b>	0.122	0.247	0.363	0.293	0.211	<i>0.063</i>	<i>0.090</i>	<i>0.066</i>
PPT × Y	0.151	0.326	0.253	0.699	0.447	<i>0.057</i>	<b>0.001</b>	<i>0.094</i>	<b>0.008</b>
C × Y	0.200	<b>0.047</b>	<b>0.021</b>	<b>0.008</b>	<b>0.010</b>	0.617	<b>0.048</b>	<b>0.012</b>	<b>0.017</b>
W × PPT × C	<b>&lt;0.001</b>	0.366	0.793	0.428	0.643	0.449	0.786	0.238	0.510
W × PPT × Y	<b>&lt;0.001</b>	<i>0.057</i>	<i>0.069</i>	<i>0.068</i>	<i>0.056</i>	0.612	<b>0.004</b>	<b>0.003</b>	<b>0.002</b>
W × C × Y	<b>&lt;0.001</b>	<b>0.021</b>	0.432	0.449	0.421	0.543	0.478	0.439	0.429
PPT × C × Y	<b>0.001</b>	0.447	0.564	0.797	0.654	0.457	0.583	0.781	0.674
W × PPT × C × Y	0.288	<b>0.041</b>	0.824	0.879	0.930	0.351	0.826	0.356	0.738

ANPP, aboveground net primary productivity; BNPP, belowground net primary productivity;  $f_{\text{BNPP}}$ , partitioning of BNPP with respect to ANPP; NPP, net primary productivity; RUE, rain-use efficiency.




**Fig. 2** Variation in aboveground net primary productivity (a), BNPP (b), NPP (c), and  $f_{\text{BNPP}}$  (d) under 12 treatments in 2010 and 2011. Values are Mean  $\pm$  SE ( $n = 6$ ). Different letters represent statistically significant differences at  $P < 0.05$ . CC, control; CD, double precipitation; CH, half precipitation; WC, warmed; WD, warmed and double precipitation; WH, warmed and half precipitation.

$\text{RUE}_{\text{BNPP}}$ , and  $\text{RUE}_{\text{NPP}}$  were positively influenced by treatment-induced changes in  $C_4/C_3$  (Fig. 6 and S1). Additionally, the interaction of altered precipitation and clipping marginally influenced  $\text{RUE}_{\text{ANPP}}$ ,  $\text{RUE}_{\text{BNPP}}$ , and  $\text{RUE}_{\text{NPP}}$  (all  $P < 0.10$ , Table 2).

*Relationships of NPP and  $f_{\text{BNPP}}$  with climatic factors and species composition*

Generally, variation in ANPP, BNPP, and NPP was positively and negatively regulated by soil moisture (all  $P < 0.01$ , Fig. 7a–c) and soil temperature (all  $P < 0.05$  except for BNPP,  $P = 0.056$ , Fig. 7e–g), respectively,




**Fig. 3** Main treatment effects of warming (a, b), double precipitation (c, d), half precipitation (e, f), and clipping (g, h) on ANPP, BNPP, NPP, and  $f_{\text{BNPP}}$  in 2010 and 2011. Values are Mean  $\pm$  SE. Asterisks indicate statistically significant difference between the control and the specific treatment at  $P < 0.05$ . ANPP, aboveground net primary productivity; BNPP, belowground net primary productivity; NPP, net primary productivity; PPT, precipitation.

under different treatments across the years. Multiple regression analysis showed that soil moisture was the dominant climatic factor that controlled the variation in ecosystem productivity. The partitioning of NPP,  $f_{\text{BNPP}}$ , was negatively correlated with soil moisture and positively correlated with soil temperature (all  $P < 0.01$ , Fig. 7d and h). Soil temperature was found to be the dominant climatic factor that regulated the variation in  $f_{\text{BNPP}}$ . Additionally, species composition, expressed as the ratio of  $C_3$  productivity to ANPP, significantly impacted ANPP, BNPP, NPP, and  $f_{\text{BNPP}}$  across the years (all  $P < 0.05$ , Fig. 7i–l).

**Discussion**


*Treatment effects on NPP*

Our results in general indicated that changes in temperature and precipitation directly and rapidly affected the




**Fig. 4** Variation in the  $RUE_{ANPP}$  (a),  $RUE_{BNPP}$  (b), and  $RUE_{NPP}$  (c) under 12 treatments in 2010 and 2011. Values are Mean  $\pm$  SE ( $n = 6$ ). Different letters represent statistically significant differences at  $P < 0.05$ . CC, control; CD, double precipitation; CH, half precipitation; WC, warmed; WD, warmed and double precipitation; WH, warmed and half precipitation; RUE, rain-use efficiency.

production of grasslands. Warming increased BNPP significantly ( $P < 0.01$ ) and NPP marginally ( $P = 0.057$ ) across the years, although its effect on ANPP was not significant ( $P > 0.05$ , Table 2; Figs 2a–c and 3a). The increases in NPP were consistent with previous studies, which have demonstrated that experimental warming that increased soil temperature ranging from 0.1 to 10.2 °C significantly increased NPP (e.g., Luo *et al.*, 2009; Wu *et al.*, 2011; Xu *et al.*, 2012). The increases in NPP may be attributed to prolonged growing seasons with earlier beginnings and later endings under continuous warming (Wan *et al.*, 2005), increased soil nutrient mineralization and plant nutrient uptake (Rustad *et al.*, 2001; Sardans *et al.*, 2008), and also stimulated photosynthetic rates at higher temperatures (Rustad *et al.*, 2001; Luo *et al.*, 2009; Wu *et al.*, 2011). However, stimulated photosynthesis may be a minor cause of the increase in NPP since warming had no significant effect on photosynthesis according to previous studies (Stirling *et al.*, 1997; Zhou *et al.*, 2007; Rogers *et al.*,



**Fig. 5** Main treatment effects of warming (a), double precipitation (b), half precipitation (c), and clipping (d) on  $RUE_{ANPP}$ ,  $RUE_{BNPP}$ , and  $RUE_{NPP}$  in 2010 and 2011. Values are Mean  $\pm$  SE. Asterisks indicate statistically significant difference between the control and the specific treatment at  $P < 0.05$ . RUE, rain-use efficiency; ANPP, aboveground net primary productivity; BNPP, belowground net primary productivity; NPP, net primary productivity; PPT, precipitation.



**Fig. 6** Relationship of the main treatment-induced changes between  $C_4/C_3$  and  $RUE_{NPPs}$  in 2010 and 2011. Treatments include warming, double precipitation, half precipitation, and clipping.  $RUE_{NPPs}$  represents  $RUE_{ANPP}$ ,  $RUE_{BNPP}$  and  $RUE_{NPP}$ .




Fig. 7 Relationships of soil moisture (%), soil temperature (°C), and species composition ( $C_3/\text{ANPP}$ ) with aboveground net primary productivity (a, e, i), BNPP (b, f, j), net primary productivity (NPP) (c, g, k), and  $f_{\text{BNPP}}$  (d, h, l) across the years ( $n = 24$ ).

2012). The non-significant warming effect on ANPP probably resulted from the translocation of photosynthates from aboveground biomass to roots under warming, which was evidenced by the increased BNPP as well as  $f_{\text{BNPP}}$  under warming (all  $P < 0.05$ , Table 2; Figs 2b, d and 3a, b).

In line with previous findings (Sala *et al.*, 1988; Knapp *et al.*, 2001; Bai *et al.*, 2008; Wu *et al.*, 2011), our results showed that altered precipitation significantly influenced ANPP, BNPP, and NPP (all  $P < 0.05$ , Table 2). Overall, double and half precipitation stimulated and suppressed plant productivity, respectively (Figs 2a–c and 3c, e). Moreover, plant productivity in 2010 was much larger than that in 2011 (all  $P < 0.01$ , Table 2; Fig. 2a–c), probably due to the precipitation amount that was 1.7 times higher in 2010 than in 2011 (Fig. 2). Water availability is crucial to terrestrial biological activities (Sala *et al.*, 1988; Li *et al.*, 2011). Observed patterns of plant productivity across natural precipitation gradients and years suggest that increases in precipitation positively stimulated plant growth in relatively dry systems (e.g., Sala *et al.*, 1988; Huxman *et al.*, 2004). At least two possible reasons may account for

clipping not significantly decreasing NPP across the years (all  $P > 0.05$ , Table 2). First, decrease in the amount of litter on the ground may increase seed bank stores and thus more germination at the beginning of growing seasons (Ruprecht & Szabo, 2012). Second, during the growing seasons, the photosynthesis of the lower canopy leaves may increase and thus give higher plant productivity due to less standing litter and improved light conditions in the clipped subplots (Xu *et al.*, 2012). Additionally, the responses of NPP to clipping varied widely by year (all  $P < 0.05$ , Table 2; Fig. 3g), probably related to the difference in precipitation amount, which in turn further confirmed the importance of precipitation in regulating plant productivity. The lack of interactive effects of warming, altered precipitation, and clipping suggest that results from single-factor manipulative experiments should be treated with caution.

#### Treatment effects on $f_{\text{BNPP}}$

The partitioning of BNPP with respect to ANPP is a critical issue in plant ecology and evolution as well as in C-cycling models (Enquist & Niklas, 2002; Ågren &

Franklin, 2003; Hui & Jackson, 2005). According to optimal partitioning theory, environmental changes may disturb the partitioning between BNPP and ANPP and cause plants to reallocate biomass among different organs to maximize their growth (Bloom *et al.*, 1985; Chapin *et al.*, 1987; Hui & Jackson, 2005). In this study, experimental warming was found to increase  $f_{\text{BNPP}}$  ( $P < 0.05$ , Table 2; Figs 2d and 3b), which has also been demonstrated in another manipulative warming experiment in a tall-grass prairie (Xu *et al.*, 2012). Moreover, in support of optimal partitioning theory, previous studies have shown that plants allocate proportionally more biomass to roots in response to warming-induced dry conditions in order to efficiently capture water (Chapin *et al.*, 1987; McCarthy & Enquist, 2007). Altered precipitation did not significantly impact  $f_{\text{BNPP}}$  across the years ( $P > 0.05$ , Table 2), probably due to the similar magnitude of precipitation-induced changes in both ANPP and BNPP, resulting in no significant net precipitation effect on  $f_{\text{BNPP}}$  (Fig. 3c–f). Clipping also had no impact on  $f_{\text{BNPP}}$  across the years ( $P > 0.05$ , Table 2) because it did not change ANPP and BNPP much (Table 2; Fig. 3g and h). Additionally,  $f_{\text{BNPP}}$  on average was smaller in 2010 than in 2011 ( $0.50 \pm 0.003$  vs.  $0.53 \pm 0.005$ ,  $P < 0.0001$ ,  $n = 48$ , Fig. 2d), indicating that low soil moisture conditions may positively impact  $f_{\text{BNPP}}$ . With global warming and changed precipitation regimes worldwide (IPCC, 2007), variation in  $f_{\text{BNPP}}$  should be taken into account in projecting climate change-terrestrial C feedback.

#### Treatment effects on RUE of NPP

Rain-use efficiency, linking the C and water cycles, is an important characteristic of ecosystem productivity and could be used to evaluate the ecosystem productivity in response to water availability (Knapp *et al.*, 2001; Huxman *et al.*, 2004; Bai *et al.*, 2008; Roupsard *et al.*, 2009). In grassland ecosystems, temperature has critical effects on RUE through its impact on evapotranspiration rates, the primary process by which precipitation is returned to the atmosphere (Wilcox *et al.*, 2006). In this study, both above- and belowground processes were examined as they responded to warming, altered precipitation, and clipping. Warming significantly increased  $\text{RUE}_{\text{BNPP}}$  but not  $\text{RUE}_{\text{ANPP}}$  and  $\text{RUE}_{\text{NPP}}$  (Table 2), suggesting that BNPP was more sensitive to warming than ANPP. In contrast to our results, warming has been reported to decrease grassland RUE as a result of reduced plant production (De Boeck *et al.*, 2006). These observations are reasonable because of differences between different grasslands in an array of factors, such as plant species composition, soil texture, and water holding capacity that influence grassland

production under warming. For example, compared with a significant warming-induced increase in ANPP in a  $C_4$  dominated grassland (Luo *et al.*, 2009), warming had no significant effect on ANPP in a  $C_3$  dominated grassland (this study).

Rain-use efficiency of NPP in response to altered precipitation was found to be very different between double precipitation and half precipitation treatments. In line with previous studies showing that irrigation decreased RUE (Huxman *et al.*, 2004; Bai *et al.*, 2008; Li *et al.*, 2011), double precipitation reduced and half precipitation increased RUE of NPP, respectively (all  $P < 0.01$ , Table 2; Figs 4 and 5b, c), confirming that water is a primary resource limiting plant growth and production (Sala *et al.*, 1988; Bai *et al.*, 2008; Xu *et al.*, 2012). Additionally, clipping marginally decreased  $\text{RUE}_{\text{ANPP}}$  and  $\text{RUE}_{\text{NPP}}$  (all  $P < 0.05$ , Table 2; Figs 4 and 5d), probably resulting from stimulated evapotranspiration due to increased soil temperature and decreased soil moisture under clipping treatment (Tables 1 and 2; Wilcox *et al.*, 2006; Xu *et al.*, 2012). Treatment-induced changes in  $C_4/C_3$  positively impacted changes in  $\text{RUE}_{\text{ANPP}}$ ,  $\text{RUE}_{\text{BNPP}}$ , and  $\text{RUE}_{\text{NPP}}$  over the years (Fig. 6 and S1), indicating a potentially essential role of plant species composition in regulating RUE and demonstrating  $C_4$  species' higher RUE than  $C_3$  species (Niu *et al.*, 2003). Moreover, the interactions of clipping with altered precipitation marginally impacted RUE of NPP (all  $P < 0.10$ , Table 2; Fig. 4), making how ecosystem RUE would change under the interactions of climate change and land use less predictable.

#### Relationships of NPP and $f_{\text{BNPP}}$ with climatic factors and species composition

Water availability, either as precipitation or soil moisture, is an overwhelmingly important controlling factor for grassland production (Sala *et al.*, 1988; Huxman *et al.*, 2004; Xu *et al.*, 2012). In accordance with previous findings that increased water availability always stimulates plant growth (e.g., Sala *et al.*, 1988; Knapp & Smith, 2001; Sherry *et al.*, 2008; Wu *et al.*, 2011), our results showed that variation of ecosystem productivity, including ANPP, BNPP, and NPP, was dominantly regulated by soil moisture across the years according to multiple regression analysis (all  $P < 0.001$ , Fig. 5a–c and e–g). Previous modeling studies also confirmed the dependence of plant productivity on soil moisture (e.g., Sherry *et al.*, 2008; Weng & Luo, 2008). Moreover, increases in N mineralization with increasing precipitation may also partly account for the increased plant production as found by Burke *et al.* (1997). The variation in  $f_{\text{BNPP}}$ , however, was mainly regulated by soil temperature ( $P < 0.01$ , Fig. 7h) based on multiple

regression analysis, although soil moisture also influenced  $f_{\text{BNPP}}$  ( $P < 0.01$ , Fig. 7d), probably because changes in temperature significantly influenced BNPP, as demonstrated by Xu *et al.* (2012).

Soil moisture, co-impacted by an array of factors such as precipitation, temperature, and land-use practices, could serve as an essential index in projecting ecosystems' responses to climatic change (Friedlingstein *et al.*, 1999; Knapp *et al.*, 2001; Sherry *et al.*, 2008). Our results showed that soil moisture increased under double precipitation and decreased under warming, half precipitation, and clipping (all  $P < 0.01$ , Table 2). The differences in soil moisture between treatments were small (Table 1), which may be due to other factors, such as precipitation distribution and soil infiltration capacity, which may also affect soil moisture (Lee *et al.*, 2007). Increases in the amount of precipitation and precipitation frequency would lead to a decrease in groundwater recharge as soil infiltration capacities are exceeded (Trenberth *et al.*, 2003; IPCC, 2007).

In addition to climatic factors, variation in plant species composition (expressed as  $C_3$  contribution to ANPP) may be another potential factor in influencing ANPP, BNPP, NPP, and  $f_{\text{BNPP}}$  (all  $P < 0.05$ , Fig. 7e–h). Our experimental site was dominated by  $C_3$  species (ca. 72% of ANPP), which played a critical role in affecting production. Previous studies have also reported the potential roles of species composition in regulating ecosystem productivity under warming, altered precipitation, and  $\text{CO}_2$  enhancement (e.g., Hoeppepner & Dukes, 2012; Polley *et al.*, 2012). The negative correlation between species composition and  $f_{\text{BNPP}}$  (Fig. 7h) suggests that dry conditions favor the growth of  $C_4$  species with an advantage in capturing water by a well-developed root system (Gao *et al.*, 2011; Bloor & Bardgett, 2012; Xu *et al.*, 2012). Moreover, plants invest more in BNPP under low water conditions (Hui & Jackson, 2005; Gao *et al.*, 2011), resulting in an increase in  $f_{\text{BNPP}}$ . However, we consider species composition to be a potential factor leading to variation in ecosystem productivity,  $f_{\text{BNPP}}$ , and RUE in this study, but our experimental design could not isolate treatment effects from species composition effects on ecosystem processes.

Our methods may underestimate overall BNPP but the estimation of treatment effects on NPP is reliable because (i) BNPP was measured once a year in this experiment while fine root turnover times could be less than a year (Gill & Jackson, 2000; Luo *et al.*, 2009); (ii) warming and clipping may not affect fine root turnover much (Fitter *et al.*, 1999; Gao *et al.*, 2008; Bai *et al.*, 2010); (iii) root turnover was stimulated by increased precipitation, leading to a further underestimation of BNPP (Bai *et al.*, 2010); and (iv) root turnover was

inhibited under potential water stress (decreased precipitation) via prolonging life span for the fine roots to main function (Norby & Jackson, 2000). Additionally, runoff could influence plant water use but it may not have had much impact on the RUE of NPP in this study because runoff is inherently a component of rainfall and our whole experimental site is very flat. In general, to accurately predict the feedback of ecosystems to climate change and land use, we have to understand the responses of ecosystem processes, such as ecosystem productivity, the partitioning of productivity, and RUE, to climate change and land use as well as their consequences for ecosystem functions.

## Acknowledgements

We would like to thank four anonymous reviewers for their comments and many laboratory members for their help with field work, especially Dr. Jianyang Xia, Dr. Lifan Jiang, and Zheng Shi. This study is financially supported by National Science Foundation (NSF) under grants DEB 0743778; and through the Midwestern Regional Center of the National Institute for Climatic Change Research at Michigan Technological University, under award number DE-FC02-06ER64158.

## References

- Ågren GI, Franklin O (2003) Root: shoot ratios, optimization and nitrogen productivity. *Annals of Botany*, **92**, 795–800.
- Bai Y, Wu J, Xing Q, Pan Q, Huang J, Yang D, Han X (2008) Primary productions and rain use efficiency across a precipitation gradient on the Mongolia Plateau. *Ecology*, **89**, 2140–2153.
- Bai W, Wan S, Niu S *et al.* (2010) Increased temperature and precipitation interact to affect root production, mortality, and turnover in a temperate steppe: implications for ecosystem C cycling. *Global Change Biology*, **16**, 1306–1316.
- Bell JE, Weng E, Luo Y (2010) Ecohydrological responses to multifactor global change in a tallgrass prairie: a modeling analysis. *Journal of Geophysical Research. Biogeosciences*, **115**, G04042, doi: 10.1029/2009JG001120.
- Bloom AJ, Chapin FS, Mooney HA (1985) Resource limitation in plants—an economic analogy. *Annual Review of Ecology and Systematics*, **16**, 363–392.
- Bloor JMG, Bardgett RD (2012) Stability of above-ground and below-ground processes to extreme drought in model grassland ecosystems: interactions with plant species diversity and soil nitrogen availability. *Perspectives in Plant Ecology, Evolution and Systematics*, **14**, 193–204.
- Brueck H, Erdle K, Gao Y, Giese M, Zhao Y, Peth S, Lin S (2010) Effects of N and water supply on water use-efficiency of a semiarid grassland in Inner Mongolia. *Plant and Soil*, **328**, 495–505.
- Burke IC, Lauenroth WK, Parton WJ (1997) Regional and temporal variation in net primary production and nitrogen mineralization in grasslands. *Ecology*, **78**, 1330–1340.
- Chapin FS, Bloom AJ, Field CB, Waring RH (1987) Plant responses to multiple environmental factors. *BioScience*, **37**, 49–57.
- Chimner RA, Welker JM, Morgan J, LeCain D, Reeder J (2010) Experimental manipulations of winter snow and summer rain influence ecosystem carbon cycling in an mixed-grass prairie, Wyoming, USA. *Ecohydrology*, **3**, 284–293.
- Cramer W, Kicklighter DW, Bondeau A *et al.*, the Participants of the Potsdam NPP Model Intercomparison (1999) Comparing global models of terrestrial net primary productivity (NPP): overview and key results. *Global Change Biology*, **5** (Suppl. 1): 1–15.
- De Boeck HJ, Lemmens CMHM, Bossuyt H *et al.* (2006) How do climate warming and plant species richness affect water use in an experimental grassland? *Plant and Soil*, **288**, 249–261.
- Dukes JS, Chiariello NR, Cleland EE *et al.* (2005) Responses of grassland production to single and multiple global environmental changes. *PLoS Biology*, **3**, 1829–1837.

- Enquist BJ, Niklas KJ (2002) Global allocation rules for patterns of biomass partitioning in seed plants. *Science*, **295**, 1517–1520.
- Epstein HE, Lauenroth WK, Burke IC, Coffin DP (1996) Ecological responses of dominant grasses along two climatic gradients in the Great Plains of the United States. *Journal of Vegetation Science*, **7**, 777–778.
- Fitter AH, Self GK, Brown TK, Bogle DS, Graves JD, Benhan D, Ineson P (1999) Root production and turnover in an upland grassland subjected to artificial soil warming respond to radiation flux and nutrients, not temperature. *Oecologia*, **120**, 575–581.
- Frank DA, McNaughton SJ (1990) Above-ground biomass estimation with the canopy intercept method – a plant growth form caveat. *Oikos*, **57**, 57–60.
- Friedlingstein P, Joel G, Field CB, Fung IY (1999) Toward an allocation scheme for global terrestrial carbon models. *Global Change Biology*, **5**, 755–770.
- Friedlingstein P, Cox P, Bets R *et al.* (2006) Climate-carbon cycle feedback analysis: results from the (CMIP)-M-4 model intercomparison. *Journal of Climate*, **19**, 3337–3353.
- Gao YZ, Giese M, Lin S, Sattelmacher B, Zhao Y, Brueck H (2008) Belowground net primary productivity and biomass allocation of a grassland in Inner Mongolia is affected by grazing intensity. *Plant and Soil*, **307**, 41–50.
- Gao YZ, Chen Q, Lin S, Giese M, Brueck H (2011) Resource manipulation effects on net primary production, biomass allocation and rain-use efficiency of two semiarid grassland sites in Inner Mongolia, China. *Oecologia*, **165**, 855–864, doi: 10.1007/s00442-010-1890-z.
- Gill RA, Jackson RB (2000) Global patterns of root turnover for terrestrial ecosystems. *New Phytologist*, **147**, 13–31.
- Gill RA, Kelly RH, Parton WJ *et al.* (2002) Using simple environmental variables to estimate below-ground productivity in grasslands. *Global Ecology and Biogeography*, **11**, 79–86.
- Groisman PY, Legates DR (1994) The accuracy of United States precipitation data. *Bulletin of the American meteorological society*, **75**, 215–227.
- Hoepfner SS, Dukes JS (2012) Interactive responses of old-field plant growth and composition to warming and precipitation. *Global Change Biology*, **18**, 1754–1768.
- Hooper DU, Johnson L (1999) Nitrogen limitation in dryland ecosystems: responses to geographical and temporal variation in precipitation. *Biogeochemistry*, **46**, 247–293.
- Hui D, Jackson RB (2005) Geographical and interannual variability in biomass partitioning in grassland ecosystems: a synthesis of field data. *New Phytologist*, **169**, 85–93.
- Huxman TE, Smith MD, Fay PA *et al.* (2004) Convergence across biomes to a common rain-use efficiency. *Nature*, **429**, 651–654.
- IPCC (2007) Summary for policymakers. In: *Climate Change 2007: The Physical Science Basis. Contribution of Working Group I to the Fourth Assessment Report of the Intergovernmental Panel on Climate Change* (eds Solomon S, Qin D, Manning M, Chen Z, Marquis M, Averyt KB, Tignor M, Miller HL), pp. 797–811. Cambridge University Press, Cambridge, United Kingdom and New York, NY, USA.
- Knapp AK, Smith MD (2001) Variation among biomass in temporal dynamics of aboveground primary production. *Science*, **291**, 481–484.
- Knapp AK, Briggs JM, Koelliker JK (2001) Frequency and extent of water limitation to primary production in a mesic temperate grassland. *Ecosystems*, **4**, 19–28.
- Lauenroth WK, Burke IC, Paruelo JM (2000) Patterns of production and precipitation-use efficiency of winter wheat and native grasslands in the Central Great Plains of the United States. *Ecosystems*, **3**, 344–351.
- Lee KS, Kim JM, Lee DR, Kim Y, Lee D (2007) Analysis of water movement through an unsaturated soil zone in Jeju Island, Korea using stable oxygen and hydrogen isotopes. *Journal of Hydrology*, **345**, 199–211.
- Li J, Lin S, Taube F, Pan Q, Dittert K (2011) Above and belowground net primary productivity of grassland influenced by supplemental water and nitrogen in Inner Mongolia. *Plant and Soil*, **340**, 253–264.
- Luo Y, Sherry R, Zhou X, Wan S (2009) Terrestrial carbon-cycle feedback to climate warming: experimental evidence on plant regulation and impacts of biofuel feedstock harvest. *Global Change Biology Bioenergy*, **1**, 62–74.
- McCarthy MC, Enquist BJ (2007) Consistency between an allometric approach and optimal partitioning theory in global patterns of plant biomass allocation. *Functional Ecology*, **21**, 713–720.
- McNaughton SJ, Banyikwa FF, McNaughton MM (1998) Root biomass and productivity in grazing ecosystem: the Serengeti. *Ecology*, **79**, 587–592.
- Meehl GA, Stocker CM, Bowker TF *et al.* (2007) Global Climate Projections. In: *Climate Change 2007: The Physical Science Basis. Contribution of Working Group I to the Fourth Assessment Report of the Intergovernmental Panel on Climate Change* (eds Solomon S, Qin D, Manning M, Chen Z, Marquis M, Averyt KB, Tignor M, Miller HL), pp. 747–846. Cambridge University Press, Cambridge, United Kingdom and New York, NY, USA.
- New M, Todd M, Hulme M, Jones P (2001) Precipitation measurements and trends in the twentieth century. *International Journal of Climatology*, **21**, 1899–1922.
- Ni J (2004) Estimating net primary productivity of grasslands from field biomass measurements in temperate northern China. *Plant Ecology*, **174**, 217–234.
- Niu S, Jiang G, Li Y, Gao L, Liu M (2003) Diurnal gas exchange and superior resources use efficiency of typical C-4 species in Hunshandak Sandland, China. *Photosynthetica*, **41**, 221–226.
- Niu S, Wu M, Han Y, Xia J, Wan S (2008) Water-mediated responses of ecosystem carbon fluxes to climatic change in a temperate steppe. *New Phytologist*, **177**, 209–219.
- Norby RJ, Jackson RB (2000) Root dynamics and global change: seeking an ecosystem perspective. *New Phytologist*, **147**, 3–12.
- Norby RJ, Luo Y (2004) Evaluating ecosystem responses to rising atmospheric CO<sub>2</sub> and global warming in a multifactor world. *New Phytologist*, **162**, 281–293.
- Polley HW, Jin VL, Fay P (2012) Feedback from plant species change amplifies CO<sub>2</sub> enhancement of grassland productivity. *Global Change Biology*, **18**, 2813–2823.
- Poorter H, Nagel O (2000) The role of biomass allocation in the growth response of plants to different levels of light, CO<sub>2</sub>, nutrients and water: a quantitative review. *Australian Journal of Plant Physiology*, **27**, 595–607.
- Rogers VL, Hoepfner SS, Daley MJ, Dukes JS (2012) Leaf-level gas exchange and foliar chemistry of common old-field species responding to warming and precipitation treatments. *International Journal of Plant Science*, **173**, 957–970.
- Roupsard O, Maire G, Nouvellon Y *et al.* (2009) Scaling-up productivity (NPP) using light or water use efficiency (LUE, WUE), from a two-layer tropical plantation. *Agroforest System*, **76**, 409–422.
- Ruprecht E, Szabo A (2012) Grass litter is a natural seed trap in long-term undisturbed grassland. *Journal of Vegetation Science*, **23**, 495–504.
- Rustad LE, Campbell JL, Marion GM *et al.*, GCTE-NEWS (2001) A meta-analysis of the response of soil respiration, net nitrogen mineralization, and aboveground plant growth to experimental ecosystem warming. *Oecologia*, **126**, 543–562.
- Sala OE, Parton WJ, Joyce LA, Lauenroth WK (1988) Primary production of the central grassland region of the United States. *Ecology*, **69**, 40–45.
- Sardans J, Peñuelas J, Prieto P, Estiarte M (2008) Drought and warming induced changes in P and K concentration and accumulation in plant biomass and soil in a Mediterranean shrubland. *Plant and Soil*, **306**, 261–271.
- Sherry RA, Weng E, Arnone JA III *et al.* (2008) Lagged effects of experimental warming and doubled precipitation on annual and seasonal aboveground biomass production in a tallgrass prairie. *Global Change Biology*, **14**, 2923–2936.
- Shipley B, Meziane D (2002) The balanced-growth hypothesis and the allometry of leaf and root biomass allocation. *Functional Ecology*, **16**, 326–331.
- Sitch S, Huntingford C, Gedney N *et al.* (2008) Evaluation of the terrestrial carbon cycle, future plant geography and climate-carbon cycle feedbacks using five Dynamic Global Vegetation Models (DGVMs). *Global Change Biology*, **14**, 2015–2039.
- Stirling CM, Davey PA, Williams TG, Long SP (1997) Acclimation of photosynthesis to elevated CO<sub>2</sub> and temperature in five British native species of contrasting functional type. *Global Change Biology*, **3**, 237–246.
- Tarr J, Botkin G, Rice EL, Carpenter E, Hart M (1980) A broad analysis of 15 sites in the tall grass prairie of Oklahoma USA. *Proceedings of the Oklahoma Academy of Science*, **60**, 39–42.
- Trenberth KE, Dai AG, Rasmussen RM, Parsons DB (2003) The changing character of precipitation. *Bulletin of the American Meteorological Society*, **84**, 1205.
- US Department of Agriculture (1972) The nation's range resources—a forest-range environmental study. USDA Forest Service Forest Resources Report 19.
- Wan S, Luo Y, Wallace LL (2002) Changes in microclimate induced by experimental warming and clipping in tallgrass prairie. *Global Change Biology*, **8**, 754–768.
- Wan S, Hui D, Wallace L, Luo Y (2005) Direct and indirect effects of experimental warming on ecosystem carbon processes in a tallgrass prairie. *Global Biogeochemical Cycles*, **19**, GB2014, doi: 10.1029/2004GB002315.
- Weng E, Luo Y (2008) Soil hydrological properties regulate grassland ecosystem responses to multifactor global change: a modeling analysis. *Journal of Geophysical Research*, **113**, G03003, doi: 10.1029/2007JG000539.
- Wilcox BP, Dowhower SL, Teague WR, Thurow TL (2006) Long-term water balance in a semiarid shrubland. *Rangeland Ecology and Management*, **59**, 600–606.
- Williams AM, Willis KE, Janes JK, Vander Schoor JK, Newton PCD, Hovenden MJ (2007) Warming and free-air CO<sub>2</sub> enrichment alter demographics in four co-occurring grassland species. *New Phytologist*, **176**, 365–374.

- Wu Z, Dijkstra P, Koch GW, Peñuelas J, Hungate BA (2011) Responses of terrestrial ecosystems to temperature and precipitation change: a meta-analysis of experimental manipulation. *Global Change Biology*, **17**, 927–942.
- Wullschlegel SD, Jackson RB, Currie WS *et al.* (2001) Below-ground processes in gap models for simulating forest response to global change. *Climate Change*, **51**, 449–473.
- Xu X, Niu S, Sherry RA, Zhou X, Zhou J, Luo Y (2012) Interannual variability in responses of belowground net primary productivity (NPP) and NPP partitioning to long-term warming and clipping in a tallgrass prairie. *Global Change Biology*, **18**, 1648–1656.
- Yahdjian L, Sala OE (2002) A rainout shelter design for intercepting different amounts of rainfall. *Oecologia*, **133**, 95–101.
- Zavaleta ES, Shaw MR, Chiariello NR, Thomas BD, Cleland EE, Field CB, Mooney HA (2003) Grassland responses to three years of elevated temperature, CO<sub>2</sub>, precipitation, and N deposition. *Ecological Monograph*, **73**, 585–604.
- Zhou X, Sherry RA, An Y, Wallace LL, Luo Y (2006) Main and interactive effects of warming, clipping, and doubled precipitation on soil CO<sub>2</sub> efflux in a grassland ecosystem. *Global Biogeochemical Cycles*, doi: 10.1029/2005GB002526.
- Zhou XH, Liu XZ, Wallace LL, Luo YQ (2007) Photosynthetic and respiratory acclimation to experimental warming for four species in a tallgrass prairie ecosystem. *Journal of Integrative Plant Biology*, **49**, 270–281.

## Supporting Information

Additional Supporting Information may be found in the online version of this article:

**Figure S1.** Relationships of the main treatment effects on  $C_4/C_3$  with changes in  $RUE_{ANPP}$ ,  $RUE_{BNPP}$ , and  $RUE_{NPP}$  under warming (a), double precipitation (b), half precipitation (c), clipping, and all four treatments (e, f). Inserted panels g, h, and i are shown to demonstrate that positive correlations exist with deleting the large values around 3 (X-axis).